

Galaxies Inside Strömgren Spheres of Luminous Quasars at $z > 6$: Detection of The First Galaxies

Renyue Cen¹

ABSTRACT

The intrinsic Ly α emission lines of normal galaxies before reionization are much absorbed by the damping wing of the Gunn-Peterson trough, rendering their direct detection nearly impossible, if their intrinsic line widths are less than ~ 100 km/s. High redshift luminous quasars prior to the completion of cosmological reionization at $z \sim 6$, on the other hand, are capable of producing large HII regions around them (Strömgren spheres) to allow their intrinsic Ly α emission lines to be transmitted without overwhelming absorption (Cen & Haiman 2000). We suggest that targeted observations at the Strömgren spheres of known luminous quasars at $z \geq 6$ would be able to detect Ly α emission lines of galaxies inside the Strömgren spheres largely unattenuated. A tunable, very narrowband filter of $\frac{\Delta\lambda}{\lambda} \sim 0.1\%$ or a narrowband filter of $\frac{\Delta\lambda}{\lambda} \sim 1\%$ with follow-up spectroscopic identifications will be required. Such observations could directly observe the sources of cosmological reionization including possibly the Pop III galaxies at $z = 6 - 20$ by JWST. Possible applications include determinations of the ionization state of the intergalactic medium, the sizes of the Strömgren spheres, the ages of the quasars, the luminosity function of high redshift galaxies and its evolution, the spatial distribution of galaxies and its evolution, the biased distribution of galaxies around quasars and the anisotropy of quasar emission. Observations using Keck-class telescopes may already be made to enable a differentiation between a fully neutral and a 10% neutral intergalactic medium at $z > 6$.

Subject headings: cosmology: theory —intergalactic medium —reionization

¹Princeton University Observatory, Princeton University, Princeton, NJ 08544; cen@astro.princeton.edu

1. Introduction

Recent observations of high redshift quasars from the Sloan Digital Sky Survey (SDSS) suggest that the cosmological reionization was completed at $z \sim 6$ (e.g., Fan et al. 2001; Becker et al. 2001, Barkana 2001; Djorgovski, Castro, & Stern 2001; Cen & McDonald 2002; Litz et al. 2002), implying a largely opaque universe beyond $z \sim 6$. In the context of the standard cold dark matter model (Bahcall et al. 1999) there is by now a clear consensus that galaxies of mass $10^8 - 10^9 M_\odot$ are largely responsible for the final cosmological reionization (e.g., Barkana & Loeb 2001; Madau 2002; Wyithe & Loeb 2003a,b; Mackey, Bromm, & Hernquist 2003; Cen 2003a,b; Venkatesan, Tumlinson, & Shull 2003; Ciardi, Ferrara, & White 2003; Somerville & Livio 2003; Fukugita & Kawasaki 2003; Sokasian et al. 2003). The recent Wilkinson Microwave Anisotropy Probe (WMAP) observations (Kogut et al. 2003) appear to confirm a much more extended reionization process, with the first reionization occurring at a redshift possibly as high as $z = 15 - 25$ (Cen 2003a). As a result, our observational frontier is quickly being pushed towards high redshifts not thought possible just a few years ago. It is of great interest to directly detect the sources of reionization and the ionization state of the intergalactic medium from $z = 6$ to $z = 30$.

To date the observed Lyman alpha emitters (LAE) at $6 < z < 6.6$ appear to be quite luminous with star formation rate (SFR) probably in excess of $40 M_\odot \text{yr}^{-1}$ (Hu et al. 2002; Kodaira et al. 2003), which, if assuming a starburst of duration of $5 \times 10^7 \text{yrs}$ and star formation efficiency of 20%, would give $\geq 1 \times 10^{10} M_\odot$ of stars. The masses of these galaxies most likely exceed $1 \times 10^{11} M_\odot$. These observed LAEs are thus much more massive than the typical galaxies thought to produce most of the ionizing photons at $z \geq 6$ – current observations are just scraping the tip of an iceberg. Because of the damping wing of the Gunn-Peterson trough cast by the significantly neutral intergalactic medium (IGM) before reionization (Miralda-Escudé 1998), the intrinsic luminosities (after correcting for possible gravitational lensing effects) for these observed LAEs may be still larger. Consistent with this, typical LAEs observed at somewhat lower redshift appear to be less luminous by a factor of ten or more (Yan, Windhorst, & Cohen 2003; Rhoads et al. 2003; Taniguchi et al. 2003). For smaller galaxies of mass $10^8 - 10^9 M_\odot$ typically with an intrinsic $\text{Ly}\alpha$ line width narrower than 100km/s, the absorption by the damping wing of intervening neutral IGM is dramatically more severe, probably rendering them undetectable (Haiman 2002).

In this *Letter* we suggest that targeted, very narrowband observations of $\text{Ly}\alpha$ emission at Strömgren spheres of known luminous quasars at $z \geq 6$ may provide a unique window to directly detect the bulk of the sources for cosmological reionization, because a large fraction of galaxies in such regions can transmit their $\text{Ly}\alpha$ emission lines largely unattenuated.

2. Galaxies Inside Strömgren Spheres of Luminous Quasars

In the absence of hydrogen recombination with a static density field, the proper radius R_S of the Strömgren sphere of a quasar of age t_Q is such that the number of hydrogen atoms in the sphere equals the total number of ionizing photons produced by the source (Cen & Haiman 2000):

$$R_S = \left[\frac{3\dot{N}_{\text{ph}}t_Q}{4\pi x_{\text{IGM}}\langle n_H \rangle} \right]^{1/3} = 4.3x_{\text{IGM}}^{-1/3} \left(\frac{\dot{N}_{\text{ph}}}{1.3 \times 10^{57} \text{s}^{-1}} \right)^{1/3} \left(\frac{t_Q}{2 \times 10^7 \text{yr}} \right)^{1/3} \left(\frac{1+z_Q}{7.28} \right)^{-1} \text{ Mpc}, \quad (1)$$

where $\langle n_H \rangle$ is the mean hydrogen density within R_S , x_{IGM} is the neutral hydrogen fraction of the IGM, and $\dot{N}_{\text{ph}}$ is emission rate of ionizing photons from the quasar. For $\langle n_H \rangle$, we have adopted the mean IGM density with $\Omega_b = 0.047$, and $\dot{N}_{\text{ph}} = 1.3 \times 10^{57} \text{s}^{-1}$ is the observed flux of SDSS 1030+0524 (Haiman & Cen 2002). In the realistic case with an evolving cosmological density field and a finite hydrogen recombination time, R_S can be computed in detail (Cen & Haiman 2000). But R_S estimated in equation (1) gives a reasonable approximation to the truth (Haiman & Cen 2002) at redshift $z \leq 8$.

At sufficiently large separations from the $\text{Ly}\alpha$ line center ($\frac{\Delta\lambda}{\lambda} \gg 2 \times 10^{-8}$) the optical depth of the damping wing is (Miralda-Escudé 1998)

$$\tau_d(\Delta\lambda) = 1.3 \times 10^{-3} x_{\text{IGM}} \frac{\Omega_b h(1-Y)}{0.03} \frac{H_0(1+z)^{3/2}}{H(z)} \left(\frac{1+z}{6} \right)^{3/2} \left(\frac{\Delta\lambda}{\lambda} \right)^{-1}, \quad (2)$$

where Ω_b is baryon density parameter, Y is the helium mass fraction of the primordial gas, H_0 and $H(z)$ are the Hubble constant at redshift zero and z , respectively, $\Delta\lambda$ and λ are the observed wavelength separation from the line center and the observed wavelength of the $\text{Ly}\alpha$ line, respectively. Replacing $\Delta\lambda/\lambda$ by $H(z)\Delta r/c$, where Δr is the distance between the surface of the Strömgren sphere and the point $\vec{r}$ inside the Strömgren sphere in question along the line of sight and c is the speed of light, and approximating $H(z) \approx H_0(1+z)^{3/2}\Omega_M^{1/2}$, we obtain

$$\tau_d(\vec{r}) = 1.2x_{\text{IGM}} \left(\frac{\Omega_M}{0.27} \right)^{-1} \left(\frac{\Omega_b}{0.047} \right) \left(\frac{\sqrt{R_S^2 - r^2 \sin^2 \theta} - r \cos \theta}{1 \text{ Mpc}} \right)^{-1}, \quad (3)$$

where r is the proper quasar-centric distance of the point in question and θ is the angle between the line segment from the quasar to the observer and the line segment from the quasar to $\vec{r}$.

The Gunn-Peterson optical depth due to the residual neutral hydrogen inside the Strömgren sphere, in the absence of other ionizing sources, can be expressed as (Cen & Haiman 2000)

$\tau_r(r) = 0.34 \left(\frac{\Omega_M}{0.27} \right)^{-1/2} \left(\frac{\Omega_b}{0.047} \right) \left(\frac{\dot{N}_{\text{ph}}}{1.3 \times 10^{57} \text{s}^{-1}} \right) \left(\frac{1+z}{7.28} \right)^{9/2} \left(\frac{r}{1 \text{ Mpc}} \right)^2$, due to a very small residual neutral fraction $x_r \leq 10^{-4}$. As a result of small x_r , the local mini-Strömgren spheres produced

by Ly α galaxies would render the absorption due to all residual neutral hydrogen inside the Strömgren sphere negligible. The total Ly α optical depth at $\vec{r}$ is hence just τ_d . From Equation (3) we see that, for a luminous quasar like SDSS 1030+0524, Ly α lines from galaxies located within the central region of $r \leq 1\text{Mpc}$, would suffer a total optical depth of $\tau_t = 0.22 - 0.36$ corresponding θ from 180° to 0° , i.e., a flux reduction of $\leq 31\%$, assuming $x_{\text{IGM}} = 1.0$. It is evident that there is a significant region within the Strömgren sphere where most of the Ly α emission galaxies are able to transmit the Ly α emission lines largely unattenuated.

There are several potentially very interesting applications from direct observations of regions around luminous quasars prior to the completion of cosmological reionization. Before turning to the possible applications it is useful to estimate the abundance of possibly detectable galaxies at $z \geq 6$. Assuming that the fraction of all mass having formed into halos with masses $M_h > 10^9 M_\odot$ is f_g and that starbursts each with a duration of t_{sb} has occurred once in each halo in the last half Hubble time, then we find that the comoving number density of such galaxies is $n_g = 1.2(\frac{t_{\text{sb}}}{5 \times 10^7 \text{yrs}})(\frac{f_g}{0.10})(\frac{M}{10^9 M_\odot})^{-1} h^3 \text{Mpc}^{-3}$. Assuming a star formation efficiency c_* we obtain the star formation rate in each galaxy $\text{SFR} = 0.68(\frac{c_*}{0.20})(\frac{t_{\text{sb}}}{5 \times 10^7 \text{yrs}})^{-1}(\frac{M}{10^9 M_\odot}) M_\odot/\text{yr}$. This simple estimate, however, involves two currently unknown parameters, c_* and t_{sb} .

Perhaps a better way to estimate the abundance and luminosity of LAE is to utilize both hierarchical CDM model and LAE survey observations at highest accessible redshift, currently at $z \sim 5.7$. Figure 1 shows the predicted cumulative number density of galaxies at $z = 5.7$ (solid curve) and $z = 6.5$ (dashed curve) for a cosmological constant dominated cold dark matter model with $\Omega_M = 0.29$, $\Lambda = 0.71$, $H = 70 \text{km/s/Mpc}$, $\sigma_8 = 0.80$ and $n = 1.0$. The observational data point for LAE shown as the solid dot in Figure 1 is taken from Rhoads et al. (2003), who give $n(\text{SFR} > 5.3 M_\odot \text{yr}^{-1}) = 5.9 \times 10^{-5} \text{Mpc}^{-3}$ at $z \sim 5.7$. We have “normalized” the observational data point to the theoretical prediction at $z = 5.7$ by requiring the same abundance, which then gives an implied mass threshold of the observed LAE by Rhoads et al. Clearly, even the LAE observed at $z \sim 5.7$ by Rhoads et al. are quite massive with masses about $10^{12} M_\odot$, if the adopted cold dark matter model is correct. The computed lower bound of star formation rate for the observed LAE at $z \sim 5.7$ by Rhoads et al. of $\text{SFR} > 5.3 M_\odot \text{yr}^{-1}$ corresponds to the flux sensitivity of $2 \times 10^{-17} \text{erg/cm}^2/\text{sec}$ of their LALA survey. If we assume that the Ly α luminosity of an LAE is simply proportional to its mass, it would require a flux sensitivity of $(10^{-18}, 10^{-19}, 10^{-20}) \text{erg/cm}^2/\text{sec}$ to detect galaxies of mass of $(10^{11}, 10^{10}, 10^9) M_\odot$ with corresponding abundance of $(0.01, 0.3, 5.0) \text{Mpc}^{-3}$ at $z = 6.5$ (open circles in Figure 1), respectively.

Between night sky lines there are a few good observational windows at $\lambda > 0.79 \mu\text{m}$ (cor-




Fig. 1.— shows the cumulative number density of galaxies as a function of mass at $z = 5.7$ (solid) and $z = 6.5$ (dashed) in a cosmological constant dominated cold dark matter model, consistent with WMAP observations, with the following parameters: $\Omega_M = 0.29$, $\Lambda = 0.71$, $H = 70\text{km/s/Mpc}$, $\sigma_8 = 0.80$ and $n = 1.0$. The solid dot is an observational data point from Rhoads et al. (2003). The open circles are three representative galaxy masses with approximate star formation rates and Ly α line fluxes indicated.

responding to $z > 5.5$ for the Ly α line), most notably the one center at $z \sim 5.7$ and the other at $z \sim 6.5$. However, even within those “good” windows the required integration time to detect fainter LAE using Keck telescopes is quite long, $t = (\text{flux}/10^{-17}\text{erg}/\text{cm}^2/\text{sec})^{-2}$ hours (Gjorgovski 2003, private communications; Fan 2003, private communications), implying that approximately a hundred hours is needed on Keck to detect galaxies of mass $10^{11} M_{\odot}$, *even in the absence of significantly attenuation by neutral IGM*. The current observational limit for Ly α flux roughly corresponds to SFR $\sim$ a few $M_{\odot}/\text{yrs}$ at $z \sim 6$ (Hu et al. 2002; Yan, Windhorst, & Cohen 2003; Rhoads et al. 2003; Taniguchi et al. 2003). Detecting even lower mass LAE may be quite infeasible using Keck telescopes. Furthermore, for ground-based observations where the primary background noise in the interested spectral regime comes from sky background, it is a very tedious task to search for usable windows between numerous strong sky lines to find high redshift galaxies and quasars. Hence, it is likely necessary to resort to James Webb Space Telescope (JWST) for detecting the bulk of galaxies (of mass $M_h \sim 10^9 M_{\odot}$) at $z > 6$. As pointed out by Oh (1999), JWST should be able to detect the H α line (and possibly H β and other Balmer lines) of galaxies of mass $M_h \sim 10^9 M_{\odot}$ at $z > 6$. A joint detection in Ly α and H α would not only firmly determine the redshift of each galaxy but also shed very useful light on important physical processes of each galaxy including dust content. In addition, the ratio of Ly α to H α lines would provide a cross-check of the damping wing modulated spatial distribution of LAE or a tool to break possible degeneracies in the spatial distribution of LAE from Ly α line alone.

We note that three factors could boost the effectiveness of star formation at high redshift in terms of Ly α emission. First, the gas at high redshift may be much metal poor; for example, taking the same Salpeter IMF would give about a factor of 2 more ionizing photons hence Ly α photons for stars with $Z \approx 10^{-2} Z_{\odot}$ compared to stars with solar metallicity. Second, if the metal poor gas at high redshift forms relatively more massive stars, the efficiency of producing ionizing photons would be higher. Third, the dust content in the high- z metal-poor galaxies should be lower, possibly allowing for relatively more Ly α photons to escape. These factors may render high redshift LAE more luminous than their low redshift counterparts at a star formation rate.

3. Applications

3.1. Probing the Ionization State of the IGM and the Size of the Strömgren Sphere

Figure (2) shows the ratio of the number of galaxies at a distance $\Delta\lambda/\lambda$ (with width $\delta\frac{\Delta\lambda}{\lambda}$) from the quasar on the blue side over the number of galaxies at the same distance from




Fig. 2.— shows the ratio of the number of galaxies at a distance $\Delta\lambda/\lambda$ (with width $\delta\frac{\Delta\lambda}{\lambda}$) from the quasar on the blue side over the number of galaxies at the same distance from the quasar on the red side within a transverse radius of $r_p = 2\text{Mpc}$ proper, as a function of $\Delta\lambda/\lambda$. Three panels correspond to three flux sensitivities of $10^{-18}\text{erg/cm}^2/\text{sec}$ (Panel a), $10^{-19}\text{erg/cm}^2/\text{sec}$ (Panel b) and $10^{-20}\text{erg/cm}^2/\text{sec}$ (Panel c), which are then translated into the threshold mass of detectable galaxies using Figure 1. In each panel two sets of cases are shown. Case (1): $R_s = 5\text{Mpc}$ proper with $x_{\text{IGM}} = (1.0, 0.1, 0.01)$, respectively, shown as solid (green), dotted (red) and dashed (blue). Case (2): $R_s = 3\text{Mpc}$ proper with $x_{\text{IGM}} = (1.0, 0.1, 0.01)$. Each set asymptotically curves downward towards the radius of the respective Strömgren sphere indicated by the vertical tick sitting on the bottom x-axis. The errorbars are statistical

the quasar on the red side within a quasar-centric transverse radius of $r_p = 2\text{Mpc}$ proper, as a function of $\Delta\lambda/\lambda$. Three panels correspond to three flux sensitivities of $10^{-18}\text{erg/cm}^2/\text{sec}$ (Panel a), $10^{-19}\text{erg/cm}^2/\text{sec}$ (Panel b) and $10^{-20}\text{erg/cm}^2/\text{sec}$ (Panel c), which are then translated into the threshold masses of detectable galaxies using Figure 1 to infer the corresponding abundances. In each panel two sets of cases are shown. Case (1): $R_s = 5\text{Mpc}$ proper with $x_{\text{IGM}} = (1.0, 0.1, 0.01)$, respectively, shown as thick solid, thick dotted and thick dashed curves. Case (2): $R_s = 3\text{Mpc}$ proper with $x_{\text{IGM}} = (1.0, 0.1, 0.01)$, respectively, shown as thin solid, thin dotted and thin dashed curves. We choose to plot the ratio instead of the direct number density to avoid ambiguity due to a possibly non-uniform, quasi spherical distribution of galaxies around quasars. Galaxies will still likely be clumpy and clustered on small scales but a large volume considered would smooth out most of the fluctuations.

Let us now estimate the accuracy of the data points. The comoving volume enclosed by a cylindrical region within a transverse radius $r_p = 2\text{Mpc}$ and a line-of-sight distance of $\Delta\lambda/\lambda = 0.01$ (from $\Delta\lambda/\lambda = -0.005$ to $\Delta\lambda/\lambda = 0.005$) is $2.1 \times 10^4\text{Mpc}^3$ comoving at $z = 6.28$, which is then multiplied by the number density of galaxies using the respective n from Figure 1 to yield the corresponding total number of galaxies above the respective mass threshold. Five data points are used for the indicated $\Delta\lambda/\lambda$ range. The errorbars are shown in each panel for the case with $R_s = 3\text{Mpc}$ proper. From Panel (a) of Figure 2 it is clear that $x_{\text{IGM}} = 1$ and $x_{\text{IGM}} = 0.1$ are distinguishable at $2 - 3\sigma$ level even for galaxies with mass $M > 10^{11}\text{M}_\odot$ (Panel a), detectable by Keck telescope with an integration time of order of 100 hours for each galaxy (assuming one spectrum per observation) with total ~ 210 galaxies. This might be a costly project but the potential payoff is attractive. Cases with $x_{\text{IGM}} = 0.1$ and $x_{\text{IGM}} = 0.01$ can only be marginally differentiated at this sensitivity level. At the same time, it is evident that the models with different R_s can be more easily distinguished, allowing for a precise measurement of R_s . In addition, a precise measurement of R_s , in conjunction with the spatial information (see §3.2), may allow a determination of the age of the quasar. With JWST, if one can reach flux levels $10^{-20} - 10^{-19}\text{erg/cm}^2/\text{sec}$, very precise measurements of x_{IGM} can be made, as can be seen from Panels (b,c) in Figure 2.

We note that $\Delta\lambda/\lambda = 0.005$ corresponds to a line-of-sight comoving distance from the quasar of 16Mpc at $z = 6.28$. The gravitational influence of the quasar host galaxy at that distance is likely to be negligible, alleviating some possible complications including infall. In addition, the typical peculiar velocities of galaxies at high redshift is likely be $< 200\text{km/s}$, which would correspond to a possible displacement of $\Delta\lambda/\lambda = 6 \times 10^{-4}$, too small to significantly matter.

Finally, we note that, in the case of a lone LAE (not sitting inside a large Strömgren

sphere of a luminous quasar), an LAE estimated to have an SFR about an order of magnitude higher than the solid dot in Figure 1 (Hu et al. 2002) can only transmit a small fraction of $\text{Ly}\alpha$ flux due to attenuation by the neutral IGM (Haiman 2002). Lower mass LAE are further devastated by their lower intrinsic line widths in addition to smaller Strömgren spheres. As Haiman (2002) pointed out, no $\text{Ly}\alpha$ line of an LAE narrower than 100km/s can be observed. Therefore, it is estimated that no LAE galaxies less massive than $10^{10} - 10^{11} M_{\odot}$ will be detectable in a random field outside a large Strömgren sphere of a luminous source.

3.2. Galaxy Luminosity Function and Spatial Distribution

Fitting the data to obtain R_s and x_{IGM} (§3.1) would yield precise information on τ_d , which in turn would allow a determination of a region on the red side of the quasar where attenuation due to intervening neutral IGM is negligible. Consequently, one can construct an unbiased luminosity function given the likely availability of up to thousands of galaxies in that region, which would further allow for corrections to be made for “missing” galaxies in other regions where τ_d is appreciable. Moreover, the spatial distribution of galaxies in this region and others in other regions would allow an accurate determination of the clustering properties of galaxies up to a scale of 10 – 30Mpc comoving.

Given the determined luminosity function and x_{IGM} , along with follow-up spectroscopic observations of the galaxies detected which would produce the line widths for these galaxies, we could self-consistently determine a fraction of these galaxies that would be directly observed even sitting in an average region and their possible observable features.

3.3. Biased Distribution Around Quasars

Taking the farthest red region as an unbiased region, this would then allow a determination of luminosity-dependent biased distributions of galaxies, down to the lowest luminosity of galaxies in the farthest red region. This may shed useful light on the formation of quasars and their host galaxies. Whether or not the farthest red region is an unbiased, “average” region can be checked using the same data, where one may expect to see a diminished radial gradient.

3.4. Probing Anisotropy of Quasar Emission

The three-dimensional information of the distribution of galaxies around a quasar makes it possible to probe the angular distribution of emission from the quasar. It should be relatively straightforward to build self-consistent models to account for possible outcomes from observations. For example, if galaxy abundance drops off rapidly at a transverse distance significantly smaller than the inferred R_s , it would indicate that the quasar emission is bipolar, whose angles can be inferred. If, in addition, there is a strong suppression of galaxies on the red side of the quasar even at small r_p , it would imply that the quasar emission is largely beamed towards us. We will not attempt to make more detailed models here. If the standard unified model for AGN/quasars (e.g., Antonucci 1993) is correct, the ensuing consequence on the distribution of LAEs around quasars will clearly be detectable. This method may provide the first and accurate way to test this picture.

In summary, we show that, with a tunable, very narrowband filter of $\Delta\lambda/\lambda \sim 0.1\%$ or a narrowband filter of $\frac{\Delta\lambda}{\lambda} \sim 1\%$ to survey a luminous quasar Strömgren sphere with follow-up spectroscopic identifications of the Ly α emission lines of individual galaxies inside the quasar Strömgren sphere, we could obtain a wealth of information for the high redshift galaxies that are responsible for reionizing the universe at $z \sim 6$. This same technique may be applicable to quasars at any high redshift. At higher redshift, galaxies are somewhat less massive, but they may have a more top-heavy IMF, possibly a higher star formation efficiency, a shorter starburst duration and much lower dust content. These positive, compensatory effects could possibly render high redshift galaxies with mass $10^7 - 10^8 M_\odot$ intrinsically as luminous as galaxies of mass $10^8 - 10^9 M_\odot$ at $z = 6$. We expect that JWST should be able to detect these galaxies at $z > 20$ (Eisenstein 2003), provided that a tunable, narrowband filter at appropriate wavelengths is available.

4. Conclusions

We show that targeted observations at the Strömgren spheres of known luminous quasars with tunable very narrowband filters of width $\sim 0.1\%$ or a narrowband filter of $\frac{\Delta\lambda}{\lambda} \sim 1\%$ to survey a luminous quasar Strömgren sphere with follow-up spectroscopic identifications of the Ly α emission lines of individual galaxies would provide a direct way to observe the Ly α emission lines of the bulk of galaxies at $z > 6$ in the mass range $10^8 - 10^9 M_\odot$ that are believed to be responsible for the second cosmological reionization. At present it appears that ground-based observations, such as by Keck, can only detect a relatively small fraction, high end tail of galaxies with mass $> 10^{11} M_\odot$. But even such observations may already be able to differentiate between a fully neutral and a 10% neutral intergalactic medium at

$z > 6$. JWST should provide the unique opportunity to detect the bulk of galaxies at high redshift. The likely gain in sensitivity with a tunable, very narrowband filter may allow JWST to detect the Pop III galaxies of mass $10^7 - 10^8 M_\odot$ with a top-heavy IMF at $z > 20$. In this case, the key is to search for luminous enough quasars at these very high redshifts, perhaps requiring an infrared survey comparable to SDSS. This is presently technologically challenging but the potential scientific payoff is immense.

We show that such observations will allow for accurate determinations of the ionization state of the IGM, the sizes of the Strömgren spheres, the ages of the quasars, the luminosity function of high redshift galaxies and its evolution, the spatial distribution of galaxies and its evolution, the biased distribution of galaxies around quasars and the anisotropy of quasar emission.

If the universe was indeed reionized twice (Cen 2003a), then there will be a window of opportunity somewhere at $z = 10 - 25$, where galaxies can transmit Ly α lines (as well as optical-UV continuum and other emission lines) unabsorbed; they should be directly observable by JWST even in the absence of luminous quasars. A signature continuum cutoff of the spectra of these galaxies shortward of $1216(1 + z_{rec,2})\text{\AA}$ would pin down the redshift of the second cosmological recombination epoch $z_{rec,2}$ and then a rapid thinning out of the number density of Ly α galaxies at some redshift $z > z_{rec,2}$ would be indicative of the first reionization epoch.

This research is supported in part by NSF grant AST-0206299. I thank George Djorgovski, Xiaohui Fan and Rogier Windhorst for useful comments and discussions, Michael Strauss for pointers to useful astronomy websites, and the referee for useful suggestions which significantly strengthen the paper.

REFERENCES

- Antonucci, R.R.J. 1993, ARAA, 31, 473
Bahcall, N.A., Ostriker, J.P., Perlmutter, S., & Steinhardt, P.J. 1999, Science, 284, 1481
Barkana, R., 2002, NewA, 7, 85
Barkana, R., & Loeb, A. 2001, Phys. Rep., 349, 125
Becker, R.H., et al. 2001, AJ, 122, 2850
Cen, R. 2003a, ApJ, 591, 12
Cen, R. 2003b, ApJ, 591, L5

- Cen, R., & Haiman, Z. 2000, *ApJ*, 542, L75
- Cen, R., & McDonald, P. 2002, *ApJ*, 570, 457
- Ciardi, B., Ferrara, A., & White, S.D.M. 2003, *astro-ph/0302451*
- Djorgovski, S.G., Castro, S., Stern, D., & Mahabal, A.A. 2001, *ApJ*, 560, L5
- Eisenstein, D. 2003, Tucson Reionization Workshop
- Fan, X., et al. 2002, *AJ*, 123, 1247
- Fukugita, M., & Kawasaki, M. 2003, *MNRAS*, 343, L25
- Haiman, Z. 2002, *ApJ*, 567, L1
- Haiman, Z., & Cen, R. 2002, *ApJ*, 578, 702
- Hu, E.M., et al. 2002, *ApJ*, 568, L75
- Kodaira, K., et al. 2003, *astro-ph/0301096*
- Litz, A., Hui, L., Zaldarriaga, M., & Scoccimarro, R. 2002, *astro-ph/0111346*, *ApJ*, in press
- Mackey, J., Bromm, V., & Hernquist, L. 2003, *ApJ*, 586, 1
- Madau, P. 2002, *astro-ph/0210268*
- Madau, P., & Rees, M.J. 2000, *ApJL*, 542, L69
- Miralda-Escudé, J. 1998, *ApJ*, 501, 15
- Oh, S.P. 1999, *ApJ*, 527, 16
- Rhoads, J.E., et al. 2003, *AJ*, 125, 1006
- Sokasian, A., Abel, T., Hernquist, L., & Springel, V. 2003, *astro-ph/0303098*
- Somerville, R.S., & Livio, M. 2003, *astro-ph/0303017*
- Taniguchi, Y., et al. 2003, *ApJ*, 585, L97
- Wyithe, S., & Loeb, A. 2003a, *ApJ*, 586, 693
- Wyithe, S., & Loeb, A. 2003b, *ApJ*, 588, L69
- Yan, H., Windhorst, R.A., & Cohen, S.H. 2003, *ApJ*, 585, L93